

## **Paleoenvironmental changes related to ice sheet and ice shelf developments in the Ross Sea since 15 kyr B.P.**

Zhihua Chen<sup>1,2</sup>, Renjie Zhao<sup>1</sup>, Shulan Ge<sup>1</sup>, Yuanhui Huang<sup>1</sup>, Zheng Tang<sup>1</sup>

<sup>1</sup>Key Laboratory of Marine Sedimentology and Environmental Geology, First Institute of Oceanography, State Oceanic Administration, Qingdao, China; chenizia@fio.org.cn

<sup>2</sup>Laboratory for Marine Geology, Qingdao National Laboratory for Marine Science and Technology, Qingdao, China.

### **Abstract**

A gravity core JB04 (172°22'21"E, 75°18'04"S, water depth 510 m), dredged from the Joides Basin in the central-western Ross Sea, was used to reconstruct paleoenvironmental changes related to ice sheet and ice shelf retreats since 15 kyr BP based on radiocarbon chronology, facies, sedimentological and geochemical analyses. A total of four facies were identified in the core from the bottom up that suggest the subglacial (314~249 cm, 15.0~11.7 kyr B.P.), sub-ice shelf (249~195 cm, 11.7~9.5 kyr B.P.), ice shelf-proximal (195~115 cm, 9.5~7.0 kyr B.P.) and modern central continental shelf environments (115~0 cm, 7.0~2.5 kyr B.P.), respectively. The grounding line retreated to the site was ca. 11.7 kyr B.P., open-marine conditions occurred at 9.5 kyr B.P. while the oceanographic and climatic settings became stable since ca. 7.0 kyr B.P. in the central-western Ross Sea.

**Keywords:** paleoenvironmental changes; ice sheet and ice shelf retreat; the Ross Sea; since 15 kyr B.P.