

Dynamic fluctuations of an East Antarctic outlet glacier since the Pliocene

Richard Selwyn Jones¹⁻³, Kevin Norton², Andrew Mackintosh^{1,2}, Jacob Anderson^{4,5}, Peter Kubik⁶, Christof Vockenhuber⁶, Hella Wittmann⁷, David Fink⁸, Gary Wilson^{4,5}, Nicholas Golledge^{1,9}, Rob McKay¹.

¹ Antarctic Research Centre, Victoria University of Wellington, PO Box 600, Wellington 6140, New Zealand;

² School of Geography, Environment and Earth Sciences, Victoria University of Wellington, PO Box 600, Wellington 6140, New Zealand;

³ Department of Geography, Durham University, Science Laboratories, South Road, Durham, DH1 3LE, United Kingdom; email: richard.s.jones@durham.ac.uk;

⁴ Geology Department, University of Otago, PO Box 56, Dunedin 9054, New Zealand;

⁵ Department of Marine Science, University of Otago, 310 Castle Street, Dunedin 9016, New Zealand;

⁶ Laboratory of Ion Beam Physics, ETH Zurich, Otto Stern Weg 5, 8093 Zurich, Switzerland;

⁷ Helmholtz-Zentrum Potsdam, GFZ Deutsches GeoForschungszentrum, Telegrafenberg, 14473 Potsdam, Germany;

⁸ Australian Nuclear Science and Technology Organization (ANSTO), Menai 2234, New South Wales, Australia;

⁹ GNS Science, Avalon, Lower Hutt, New Zealand;

Abstract

Understanding past changes in the Antarctic ice sheets provides insight into how they might respond to future climate warming. The degree to which Antarctic ice sheets expanded and contracted under past climates remains uncertain, especially prior to the Last Glacial Maximum. During Pliocene and Pleistocene glacial-interglacial cycles, the East Antarctic Ice Sheet remained relatively stable in its interior (e.g. Yamane et al., 2015) and oscillated at its marine-based margin (e.g. Naish et al., 2009; Pollard & DeConto, 2009). It is currently not clear how outlet glaciers, which connect the ice sheet interior to its margin, responded to these orbitally-paced climate cycles.

We show new ice surface constraints from Skelton Glacier, an outlet of the East Antarctic ice sheet, which drains into the Ross Ice Shelf and over the AND-1B drill site (Talarico et al., 2012). Our multiple-isotope (¹⁰Be and ²⁶Al) data indicate that currently ice-free areas adjacent to the glacier underwent substantial periods of ice cover and exposure in the past. We determine the probable ice surface history implied by our data by adopting a burial-exposure modelling approach (Bierman et al., 1999), driven by orbitally-paced oscillations of the ice margin (as recorded at AND-1B; Naish et al., 2009) over the period from the Pliocene to the present day. This analysis shows that: 1) the glacier surface has likely fluctuated since at least the Pliocene; 2) the ice surface was >200 m higher than today during glacial periods, and the glacier has been thicker than present for ~75–90% of each glacial-interglacial cycle; and 3) ice cover at higher elevations possibly occurred for a relatively shorter time per Pliocene cycle than Pleistocene cycle. While it is hard to determine the magnitude of surface lowering during interglacial periods, our data shows that the duration of deglaciation, relative to glacial expansion, was similar in both the Pliocene and Pleistocene.

These outcomes are consistent with evidence from the edge of the East Antarctic plateau that indicate a higher ice sheet surface existed during the Pliocene (e.g. Balco et al., 2014; Di Nicola et al., 2012; Yamane et al., 2015), as well as other data collected from sites at lower altitudes and nearer to the coast, suggesting that glacier surface elevations fluctuated substantially since the Pliocene (e.g. Balco et al., 2014; Di Nicola et al., 2009; Lilly et al., 2010). Large glacial-interglacial fluctuations in ice extent occurred at the marine-based ice margin of the East Antarctic Ice Sheet, in response to the evolving climate and bed topography since at least the Pliocene (e.g. Levy et al., 2012, Pollard & DeConto, 2009). Over this time, our data and analysis demonstrate that such dynamic changes extended upstream to outlet glaciers that connect the ice sheet interior to the ice margin.

Keywords: cosmogenic nuclides, Pliocene, Pleistocene, EAIS

References

- Balco, G., Stone, J.O.H., Sliwinski, M.G., Todd, C., 2014. Features of the glacial history of the Transantarctic Mountains inferred from cosmogenic ^{26}Al , ^{10}Be and ^{21}Ne concentrations in bedrock surfaces. *Antarctic Science* 26, 708-723.
- Bierman, P.R., Marsella, K.A., Patterson, C., Davis, P.T., Caffee, M., 1999. Mid-Pleistocene cosmogenic minimum-age limits for pre-Wisconsinan glacial surfaces in southwestern Minnesota and southern Baffin Island: a multiple nuclide approach. *Geomorphology* 27, 25-39.
- Di Nicola, L., Baroni, C., Strasky, S., Salvatore, M.C., Schlüchter, C., Akçar, N., Kubik, P.W., Wieler, R., 2012. Multiple cosmogenic nuclides document the stability of the East Antarctic Ice Sheet in northern Victoria Land since the Late Miocene (5–7 Ma). *Quaternary Science Reviews* 57, 85-94.
- Di Nicola, L., Strasky, S., Schlüchter, C., Salvatore, M.C., Akçar, N., Kubik, P.W., Christl, M., Kasper, H.U., Wieler, R., Baroni, C., 2009. Multiple cosmogenic nuclides document complex Pleistocene exposure history of glacial drifts in Terra Nova Bay (northern Victoria Land, Antarctica). *Quaternary Research* 71, 83-92.
- Levy, R., Cody, R., Crampton, J., Fielding, C., Golledge, N., Harwood, D., Henrys, S., McKay, R., Naish, T., Ohneiser, C., Wilson, G., Wilson, T., Winter, D., 2012. Late Neogene climate and glacial history of the Southern Victoria Land coast from integrated drill core, seismic and outcrop data. *Global and Planetary Change* 96-97, 157-180.
- Lilly, K., Fink, D., Fabel, D., Lambeck, K., 2010. Pleistocene dynamics of the interior East Antarctic ice sheet. *Geology* 38, 703-706.
- Naish, T., Powell, R., Levy, R., Wilson, G., Scherer, R., Talarico, F., ... 2009. Obliquity-paced Pliocene West Antarctic ice sheet oscillations. *Nature* 458, 322-328.
- Pollard, D., DeConto, R.M., 2009. Modelling West Antarctic ice sheet growth and collapse through the past five million years. *Nature* 458, 329-332.
- Talarico, F.M., McKay, R.M., Powell, R.D., Sandroni, S., Naish, T., 2012. Late Cenozoic oscillations of Antarctic ice sheets revealed by provenance of basement clasts and grain detrital modes in ANDRILL core AND-1B. *Global and Planetary Change* 96-97, 23-40.
- Yamane, M., Yokoyama, Y., Abe-Ouchi, A., Obrochta, S., Saito, F., Moriwaki, K., Matsuzaki, H., 2015. Exposure age and ice-sheet model constraints on Pliocene East Antarctic ice sheet dynamics. *Nature Communications* 6, 7016.

