

The role of the ocean for Antarctic cryosphere dynamics during mid Miocene: a view from offshore Wilkes Land

Francesca Sangiorgi^{1*}, Peter K. Bijl¹, Sandra Passchier², Ulrich Salzmann³, Stefan Schouten^{4,5}, Robert McKay⁶, Rosie D. Cody^{6#}, Jörg Pross⁷, Tina van de Flierdt⁸, Steven M. Bohaty⁹, Richard Levy¹⁰, Trevor Williams¹¹, Carlota Escutia¹², Henk Brinkhuis^{1,4}

¹ Marine Palynology and Paleoceanography, Laboratory of Palaeobotany and Palynology, Department of Earth Sciences, Faculty of Geoscience, Utrecht University, Heidelberglaan 2, 3584CS Utrecht, The Netherlands; f.sangiorgi@uu.nl

² Department of Earth and Environmental Studies, Montclair State University, 1 Normal Ave, Montclair, New Jersey, U.S.A.;

³ Department of Geography, Faculty of Engineering and Environment, Northumbria University, Newcastle upon Tyne, NE1 8ST, UK;

⁴ NIOZ Royal Netherlands Institute of Sea Research, Texel, The Netherlands;

⁵ Organic Geochemistry, Department of Earth Sciences, Faculty of Geoscience, Utrecht University;

⁶ Antarctic Research Centre, Victoria University of Wellington, PO Box 600, Wellington, 6140, New Zealand;

⁷ Paleoenvironmental Dynamics Group, Institute of Earth Sciences, Heidelberg University, Im Neuenheimer Feld 234, 69120 Heidelberg, Germany;

⁸ Imperial College London, Department of Earth Science and Engineering, South Kensington Campus, Exhibition Road, London, SW7 2AZ, UK;

⁹ Ocean and Earth Science, University of Southampton, National Oceanography Centre Southampton, European Way, Southampton SO14 3ZH, UK;

¹⁰ Department of Paleontology, GNS Science, 1 Fairway Drive, Lower Hutt 5040, New Zealand

¹¹ International Ocean Discovery Program, Texas A&M University, 1000 Discovery Drive, College Station, TX 77845, USA;

¹² Instituto Andaluz de Ciencias de la Tierra, CSIC-University of Granada Av. de las Palmeras, 4, 18100 Armilla (Granada), Spain.

Abstract

With the ongoing increase in atmospheric CO₂ and global temperatures, a fundamental scientific and societal question arises concerning the stability of the Antarctic cryosphere. Modern observational data indicate the Southern Ocean has experienced significant warming, with oceanic fronts being pushed several tenth of km closer to the continent. Moreover, basal melt of ice shelves from warming oceans is causing accelerated grounding line retreat of the Antarctic ice sheets and shelves. However, monitoring data are available for the last few decades only, which prevents the evaluation of long-term changes in ice mass balance. Studying intervals in Earth's past history, which represent the best possible analogues of (near) future conditions, becomes thus essential.

The Miocene Epoch encompasses periods with CO₂ concentrations between today's and those expected for the (near) future. It has also become clear that ice-proximal oceanographic regime is a critical factor for the stability and mass balance of ice sheets.

We investigate the role of the ocean in the (in)stability of the Wilkes Land subglacial basin (East Antarctica) during the climatically highly variable mid-Miocene between ~17-11 million years [Ma] ago (albeit with a hiatus at around 13.4 Ma) by studying marine sediment cores from offshore Adélie Coast, retrieved during Integrated Ocean Drilling Program (IODP) Expedition 318 at Site U1356. Warm surface-ocean and continental temperatures, absence of sea-ice, and abundant soil formation and erosion along the coast imply that the East Antarctic ice sheet was reduced further than its terrestrial margin during the mid-Miocene Climatic Optimum (MCO; 17–15 Ma). After the MCO, a marine-based ice sheet developed, though sensitive to melting upon contact with warm ocean waters. Comparing these data with available reconstructions from

the Ross Sea (ANDRILL, AND-2A record) and the ODP Site 1171 (south of Tasmania), our results suggest that the mid-Miocene temperature gradient never resembled that of present-day and demonstrate the importance of oceanic climate on Antarctic continental conditions. This study confirms present-day observations of Eastern Antarctic sub-glacial basins highly sensitive to ocean warming.

Keywords: Miocene, Wilkes Land, palynology, ocean temperature, vegetation