

## Sea ice and diatom productivity in the Ross Sea, Antarctica: the response to post-glacial warming

Rebecca Parker<sup>1</sup>, Christina Riesselman<sup>1,2</sup>, Jae Il Lee<sup>3</sup>, Min Kyung Lee<sup>3</sup>, Geraldine Jacobsen<sup>4</sup>, Brad Rosenheim<sup>5</sup>, Cristina Subt<sup>5</sup>, Atun Zawadzki<sup>4</sup>, Robert McKay<sup>6</sup>, Richard Levy<sup>7</sup>, Kyu-Cheul Yoo<sup>3</sup>

<sup>1</sup>Department of Geology, University of Otago, Dunedin 9054, New Zealand;

[rebeccaparkernz@gmail.com](mailto:rebeccaparkernz@gmail.com)

<sup>2</sup>Department of Marine Science, University of Otago, Dunedin 9054, New Zealand;

<sup>3</sup>Korean Polar Research Institute (KOPRI), Incheon 406-840, South Korea;

<sup>4</sup>Australian Nuclear Science and Technology Organisation (ANSTO), Lucas Heights, NSW, Australia;

<sup>5</sup>College of Marine Science, University of South Florida, St. Petersburg, FL, USA;

<sup>6</sup>Victoria University, Wellington, New Zealand;

<sup>7</sup>GNS Science, Lower Hutt, New Zealand.

### **Abstract**

The Holocene is characterised by millennial-scale climate fluctuations superimposed on an orbitally-paced warming trend. Spatial heterogeneity in these fluctuations appears to result from local variations in climate and oceanography associated with deglaciation. While timing of ice-retreat within the Ross Sea has been regionally constrained (Jones et al., 2015; McKay et al., 2016), there are fewer constraints on oceanographic and climatic conditions during Holocene warming, and the subsequent impacts on sea-ice extent and primary productivity.

We present a new multi-site reconstruction of deglacial ice-retreat timing and Holocene sea-ice extent and primary production from four Holocene marine sedimentary records. These cores were collected from the southwestern Ross Sea during the 2015 *Araon* Cruise ANA05B, and are located offshore of Mawson Glacier (GC78), Granite Harbour (GC80), north and northeast of Ross Island (GC72 and GC71 respectively). Cores are up to 4.5 m long and contain basal ice-associated facies overlain by diatom ooze. Core-to-core correlation is achieved through ramped pyrolysis <sup>14</sup>C chronology, reservoir-corrected using <sup>210</sup>Pb dating (Rosenheim, 2013). It is a unique chronological approach to this region, whereby local reservoir ages have been generated. Comparison to published records from onshore and offshore indicates a complex pattern of ice-shelf retreat in this region, and helps refine the history of Holocene variability in the SW Ross Sea.

To reconstruct the post-glacial environment, we utilize geochemical proxies for gross diatom abundance, nutrient utilization and water stratification (wt.% BSi,  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ), and diatom assemblages for sea-ice extent. Of particular interest is a  $\delta^{13}\text{C}$  excursion ( $\sim$ -30‰) seen in GC72, GC78 and GC80 at 65, 95 and 104 cmbsf respectively. This excursion corresponds to a high abundance of *Corethron* spp. and/or *Chaetoceros* spp. that reflects a transient change in oceanography and sea ice dynamics. Similar monospecific *Corethron* spp. layers have been observed in several regions around the Antarctic continent that reflect a transient environment controlled by unusually warm conditions (Leventer et al., 1993; Taylor and McMin, 2001; Maddison et al., 2011).

**Keywords:** paleoclimate, diatom micropaleontology, isotope geochemistry, radiocarbon

## **References**

- Jones R.S., Mackintosh A.N., Norton, K.P., Golledge, N.R., Fogwill, C.J., Kubik, P.W., Christl, M., Greenwood, S.L. 2015. Rapid Holocene thinning of an East Antarctic outlet glacier driven by marine ice sheet instability. *Nat. Comms*, 6 1-9.
- Leventer, A., Dunbar, R.B., DeMaster, D.J. 1993. Diatom Evidence for Late Holocene Climatic Events in Granite Harbour, Antarctica. *Paleoceanography*, 8(3) 373-386
- Maddison, E.J., Pike, J., Dunbar, R. 2011. Seasonally laminated diatom-rich sediments from Dumont d'Urville Trough, East Antarctic Margin: Late-Holocene Neoglacial sea-ice conditions. *The Holocene*, 22(8) 857-875
- McKay, R.M., Golledge, N.R., Maas, S., Naish, T., Levy, R., Dunbar, G. and Kuhn, G., 2016. Antarctic marine ice-sheet retreat in the Ross Sea during the early Holocene. *Geology*, 44(1), 7-10
- Rosenheim, B.E., Santoro, J.A., Gunter, M., and Domack, E., 2013. Improving Antarctic Sediment <sup>14</sup>C Dating Using Ramped Pyrolysis: An Example from the Hugo Island Trough. *Radiocarbon*, 55(1) 115-126
- Taylor, F and McMinn, A. 2011. Evidence from diatoms for Holocene climate fluctuations along the East Antarctic margin. *The Holocene*, 11(4) 455-466